

Education information



Working together with individuals, families and medical professionals to offer support and information on syncope and reflex anoxic seizures

The aims of this booklet

- ★ To inform and reassure you.
- ★ To outline the ways a school can effectively manage a child with syncope/RAS so as to provide support and enable inclusion.
- ★ To provide you with the information, materials and support with which you can plan and then action a whole school strategy on the care and management of a child with syncope/RAS.



Definitions

- ★ It is believed that about two thirds of loss of consciousness is due to syncope. Syncope (sin-co-pee) is a transient loss of consciousness and posture, caused by a temporary reduction in blood flow to the brain. It is characterised by rapid onset, short duration, and spontaneous complete recovery.
- ★ This is quite different from epilepsy, where the problem begins in the brain itself, and the blood flow during an attack may be completely normal.
- ★ Reflex Anoxic Seizures (RAS) is the childhood version of Reflex Syncope.
- ★ Reflex Syncope is similar to RAS, but is more commonly diagnosed in older patients.

Key points to remember

This is a NON-life threatening condition and there are no known long term ill effects.



- ♥ The frequency of attacks varies with each individual – they may occur several times per day/ week/ month or year.
- ♥ While RAS more frequently occurs in young children, it is not uncommon for it to occur at any age.
- ♥ Good management of the child is the key to inclusion.

What will I see when someone is having an attack?

Each child is individual so symptoms may differ slightly

Triggers

Unexpected stimulus – such as pain, shock, surprise or fright

Signs

- May be off colour
- A marked pallor of the skin
- Emotional
- Keeping a diary can help predict episodes

Symptoms

The heart and breathing momentarily stop. The eyes roll up into the head, the complexion becomes deathly white, (while the skin may appear blue around the mouth and under the eyes), and the jaw clenches and the body stiffens. There may be loss of muscle tone and possibly loss of bladder control

What will I see when someone is entering the recovery stage?

Recovery

The body relaxes and the heart starts beating (sometimes very slowly). Consciousness is usually regained within one or two minutes, but occasionally it may be longer.

Upon Recovery

The child may be very emotional and fall into a deep sleep for two or three hours. They may also look extremely pale with dark circles under their eyes.

Think of a hedgehog! Whenever the hedgehog faces danger it curls into a ball shutting itself off from the world to protect itself and only uncurls when it is sure it is safe to do so. This is very similar to children with RAS, their body shuts down until the 'nasty thing' has gone away.

What should I do if someone is having an attack?

Don't panic

An unexpected syncopal attack can be alarming, just remember that the child WILL recover.

Roll the child into recovery position.

Are there any dangers which could cause them harm? If so, make the area safe.

Stay with them, talking quietly and reassuringly. To the recovering child everything can appear disorientated and noisy, so try to keep the room calm, quiet and be reassuring!

The Recovery Position

1. Place arm nearest to you at a right angle.
2. Move other arm, and place the back of their hand against their opposite cheek. Then get hold of the knee furthest from you and pull up until foot is flat on the floor.



3. Pull the knee towards you, keeping the child's hand pressed against their cheek, and position the leg at a right angle.
4. Make sure that the airway remains open by tilting the head back and lifting the chin, and checking the breathing.
5. Monitor the child's condition until help arrives, if help is needed.

How can I help to prevent attacks?

- ♥ Share information with ALL staff who come into contact with the child.
- ♥ Keep a note of days of poor concentration, it can be a forewarning of possible episodes.
- ♥ Maintain good hydration – drink lots of water.
- ♥ Pre-warn the child of possible dangers, to reduce unexpected shock.
- ♥ Allow the child to watch, survey and become accustomed before participating in any activity whether that be a game, party or even a crowded room.
- ♥ Swimming: encourage them to participate, but allow them time to enter the water gradually.

During and after an attack

Stay calm!

- Make sure the child is safe. Lie the patient down flat – on your lap / floor / bed or sofa and put them into the recovery position. Check their airway to make sure nothing is in their mouth.
- When the child regains consciousness allow them to rest, and only move them when they feel ready. Talk to them in a quiet voice constantly reassuring them, even give them a cuddle.
- If they wish to sleep, allow it (they may sleep for two-three hours). Or they may feel dizzy, afraid and disorientated so try to reassure them.

- Many report that noise is amplified, so try to keep the environment around them as quiet and calm as possible.
- The child may be emotional for quite a while after the event – on an individual basis, decide whether they need to be sent home.
- If an injury is sustained, or you are concerned, or the child is unconscious for an unusually long period, please call the emergency services.
- If they are not breathing or have no pulse dial 999 immediately and start basic life support.

How do I manage behaviour of a pupil with RAS?

Knowing that unexpected bumps and shocks can trigger an attack, it is understandable that some parents and teachers alike find managing behaviour worrying. However it should be remembered that:

- ♥ Children with RAS or syncope have a medical condition but should be expected to behave as any other pupil.
- ♥ Occasionally a loud, unexpected shout or noise could trigger an attack, such as the sound of the fire alarm or being startled by a shout.
- ♥ Anxiety can also trigger 'dizzy' spells.
- ♥ After an attack it is best if they can recover in a quiet area. For some it may be necessary to be sent home. However, once they have recovered and returned to the class they should be able to return to normal class activities and follow usual codes of conduct.
- ♥ If uncertain, seek advice from the family or contact www.stars.org.uk or 01789 867 503 or info@stars.org.uk.

How can I enable inclusion?

- ♥ Share the child's care plan with all staff who will be responsible for the care of the pupil.
- ♥ Maintain regular communication with the child's family.
- ♥ Educate other students about RAS and syncope.
- ♥ Plan ahead time to share information with the pupil - for example when an alarm is due to ring.
- ♥ Allow time to acclimatise - to gently slide into a swimming pool or watch an activity before joining in.
- ♥ If uncertain, seek advice from the family or contact www.stars.org.uk or 01789 867 503 or info@stars.org.uk.



What should be included in a care plan?

Download a care plan from the STARS website: www.stars.org.uk.

- ♥ Student name
- ♥ Student DOB
- ♥ Student photo
- ♥ Background of medical condition
- ♥ Triggers & warning signs
- ♥ Management strategy

Important Information

Emergency Contact Details:

Parent/Guardian

Doctor

STARS helpline: 01789 867 503

Other information

Signed & Dated by:

Head Teacher

Parent/Guardian

Student

Frequently Asked Questions

What is Reflex Anoxic Seizures?

RAS occur mainly in young children but can occur at any age. Any unexpected stimulus, such as pain, fright or shock, causes the heart and breathing to stop. The eyes roll up into the head, and the complexion becomes deathly white, often blue around the mouth and under the eyes. Also the jaw may clench and the body stiffen. Sometimes the arms and legs will jerk. After about 30 seconds the body relaxes and the heart and breathing restart, but the sufferer remains unconscious. One or two minutes later the person may regain consciousness but can appear to be unconscious for over an hour.

Upon recovery the person may be emotional and sleep for three to four hours, RAS can occur several times a day/ week/month and often occur in clusters.

What is the age of the youngest and oldest known RAS sufferer?

The youngest known sufferer had his first attack during his first day of life. There are RAS sufferers in their 50s and 60s.

What causes RAS in children?

Reflex Anoxic Seizures are caused by the vagus nerve being over stimulated due to an unexpected stimulus such as pain/shock/fear. It is not known why this should happen in some children.

Will the heart weaken through time?

No, an RAS attack puts no strain at all on the heart.

Placing the child into the recovery position immediately after the fright or other stimuli and before loss of consciousness seems to lessen the severity of the attack, or result in a 'near miss'. Is there any medical foundation for this?

This does seem to be the case, although the doctors are unsure why. It should be noted that the child does not always have to be placed in the recovery position. (Possibly it reduces the effect of the drop in blood pressure which occurs after the heart stops.)

Does an RAS attack cause any damage to the brain, short term/long term?

There is no evidence that RAS causes brain damage. The 'fail safe' mechanism in the brain restarts the heartbeat and breathing before the oxygen level is low enough to cause damage.

If the cessation of the heart beat and breathing continues beyond the child's normal reaction time at what point should resuscitation be attempted, and how should this be administered?

It would be extremely unlikely that resuscitation would be needed because of the 'fail safe' mechanism in the brain. However, if the child's heart and breathing has not restarted after a period of two minutes, then attempting resuscitation in the normal manner would do no harm.

Is there a support group for them?

Yes. STARS, Syncope Trust And Reflex anoxic Seizures, provide support for sufferers of all ages.

Can an RAS attack occur while the child is sleeping?

Probably not, however there are other heart stopping conditions, notably Prolonged QT Syndrome, which do occur during sleep, and while awake especially with exercise. Prolonged QT Syndrome can be diagnosed using an electrocardiogram (ECG), and parents are encouraged to have this done, to rule out this diagnosis.

Is there a genetic factor in RAS?

Almost certainly. More research is needed to establish what gene actually carries the susceptibility to RAS.

Could the child with RAS carry a card like people with epilepsy or diabetes, with 'RAS' on one side and instructions for first aid response on the other?

Medic Alert pendants or wrist tags are appropriate for a child with RAS. STARS also offer an information card with the child's name on one side and medical instructions on the other. This could be kept in a purse, wallet, bag or pocket. For further information on alert cards contact info@stars.org.uk.

Is there a link between RAS and other medical conditions, e.g. early childhood illness, heart murmurs or developmental delays?

As far as the medical experts know, the answer is no.

What can I say to a child to comfort them during an attack?

It does seem to help to talk to your child calmly and reassuringly during an attack as it would appear that they are aware of at least the sound of voices. Perhaps talk about the child's favourite meal, holiday, activities, birthday parties etc.

Has an RAS patient had an attack in a swimming pool? If so, what happened?

Yes - fortunately the parent noticed immediately so no harm occurred. The stimulus of cold water splashing in the face, is a particularly strong stimulus to the vagus nerve, and can result in an attack. The child with RAS must learn to swim but should have close parental supervision at all times in the pool.

Are pains in the legs common amongst children with RAS?

Yes, however many children suffer with pains in their legs even if they do not have RAS .

What is the longest time a child with RAS has been free from attacks before they experience attacks again?

10 years.

Why does a child's face in an RAS attack go white and not blue?

During an RAS attack the heart stops beating, and breathing stops. Blood vessels constrict and the blood is not able to get to the skin. This results in white pallor to the face. The blood is lacking in oxygen and will be blue coloured; only noticeable on the lips in the early stages of the attack. During a blue breath-holding attack lack of oxygen from the lungs causes blood to turn blue. However, as the heart is still pumping, this 'blue' blood passes through the blood vessels of the face, causing the whole face and the skin of the body to have a blue tinge.

What makes the heart restart after an attack?

As the oxygen level decreases in the brain, there is a chemical release (the fail-safe mechanism), which induces the heartbeat and breathing mechanisms to restart.

How does an epileptic attack differ from an RAS attack?

An EEG (brain wave activity) of an epileptic attack would show excessive elective spike or epileptic discharges, which are not present in an RAS attack. In an RAS attack the EEG goes slow for a few seconds then is flat as the cerebral cortex shuts down, then goes slow again during the drowsy recovery phase before returning to normal. An RAS attack is always precipitated by a shock trigger, albeit sometimes difficult to identify.


Are there any problems with insurance cover for RAS students?

Schools, playgroups etc, should be made aware of the child's condition, and should advise their insurance companies accordingly. Travel insurance companies should be informed of the child's condition and may require a doctor's letter to certify that the person is fit to travel. This should not increase your premium.

Is there a pattern to the frequency of the attacks?

It does appear that they occur in batches, as yet, we don't know why.

Syncope Education DVD for schools and colleges



A Syncope Education DVD has been produced to provide information for staff on how to care for young people diagnosed with syncopal conditions. To obtain this DVD contact STARS.

For further information, help and support please contact: www.stars.org.uk
info@stars.org.uk

Telephone: +44 (0) 1789 867 503

STARS, PO Box 175, Stratford upon Avon, CV37 8YD

Donation Form **STARS**

Syncope Trust And Reflex anoxic Seizures®

Your details

Title _____ First Name(s) _____ Surname _____

Address _____

Postcode _____

Email _____ Phone number _____

Your donation Please tick method of payment & complete relevant section:

enclose a cheque made payable to 'STARS' for £ _____

or Please debit my credit/debit card for £ _____ Card type maestro visa mastercard
Valid from / Card no.
Expiry date / 3-digit Security no.

or I would like to set up a regular standing order for £ _____ starting on / /
To be paid: Monthly Quarterly Annually
My bank name _____ Bank address _____

Account no.
Sort code

Payable to: Syncope Trust And Reflex anoxic Seizures, Account no. 02455170, Sort code. 30-98-26
Lloyds TSB Plc, 22 Bridge Street, Stratford upon Avon, CV37 6AG
Please note you can cancel your standing order at any time by contacting your bank.

Gift Aid

If you pay UK Tax, the Government will give us 25% on top of your donation at no cost to you.
In order to Gift Aid your donation you must tick the box below:

* I want to Gift Aid my donation of £ _____ and any donations I make in the future
or have made in the past 4 years to STARS.

*I am a UK taxpayer and understand that if I pay less Income Tax and/or Capital Gains Tax than the amount of Gift Aid claimed on all my donations in that tax year it is my responsibility to pay any difference. Please notify us if you want to cancel this declaration, change your name or home address or no longer pay sufficient tax on your income and/or capital gains. Your address is needed to identify you as a current UK taxpayer.

Your Signature

Signature _____

Date / /

Post me to: STARS, PO Box 175, Stratford upon Avon, CV37 8YD

If you have any queries please do not hesitate to call us on 01789 867503

Registered charity number 1084898

STARS

Syncope Trust And Reflex anoxic Seizures®

Working together with individuals, families and medical professionals to offer support and information on syncope and reflex anoxic seizures

Please remember that this publication provides general guidelines only. Individuals should always discuss their condition with a healthcare professional.

STARS would like to thank all those who helped in the development and review of this publication. In particular, thanks are given to Dr William Whitehouse and Dr Charlotte D'Souza.

Unit 6B
Essex House
Cromwell Business Park
Chipping Norton
Oxfordshire
OX7 5SR

+44(0)1789 867 503

@ info@stars.org.uk

➔ www.stars.org.uk

Registered Charity
No. 1084898

© STARS
Published July 2010,
Reviewed July 2017



endorsed by
 Department
of Health

Patrons

Prof. J B P Stephenson

Dr A P Fitzpatrick

Prof. R A Kenny

Prof. C J Mathias

Dr W Whitehouse

Dr W Wieling

Dr P Boon Lim

Sir Elton John

Mr John Burton Race

Mrs Twiggy Lawson

Selma Blair

Trustees

Mr Andrew Fear

Mrs Cathrine Reid

Prof. Rose Anne Kenny

Dr William Whitehouse

Founder & Chief Executive

Trudie Lobban MBE

If you would like further information or would like to provide feedback please contact STARS.

